

Physics 161: Homework 3

(19 January 2000, due 26 January)

Problems

1. Consider the [power spectrum](#) for the van der Pohl oscillator driven at frequency 1.15 available under the “[Homework](#)” link on the website. We might be interested in the question of whether the motion is periodic, i.e. locked to the drive frequency, or quasiperiodic, i.e. the internal oscillations at a frequency $\omega_{in} \simeq 1$ and the drive at a frequency $\omega_D = 1.15$ both present (together with other frequencies, $m\omega_{in} \pm n\omega_D$ with m, n integers, given by the non-linear interaction).
 - (a) With the values of *dt*, *Interval*, and *Points* initially set for the applet would you expect to be able to resolve this question and why? (You might want to read the [instructions](#) and discussion of the [diagnostics](#) for the *Odes* applet on the website.)
 - (b) Change the values of any of *dt*, *Interval*, *Points*, and *Window* appropriately to get the best information you can on the spectrum, and make a sketch of what you see.
 - (c) Is the motion probably periodic or quasiperiodic?

2. The Hénon map is given by:

$$\begin{aligned}x_{n+1} &= 1 - ax_n^2 + y_n \\y_{n+1} &= bx_n\end{aligned}$$

Common parameters used are $a = 1.4$ and $b = 0.3$.

- (a) Set up a scheme for calculating the two Lyapunov eigenvalues of the map, i.e. do all the algebra to make explicit what you would iterate and how you would calculate the exponents.
 - (b) Use the [2dmap](#) demonstration to study the variation of the largest Lyapunov exponent on the Hénon attractor with a between 0.2 and 1.4 for $b = 0.3$ and relate the variation to the nature of the attractor. (Note: make sure you hit the *Reset* button on the applet after you change a to restart the calculation of the Lyapunov exponent. For the periodic orbits with a few points only you might want to increase *Mark* to 0.5 to make the points more obvious.)
3. Consider the two dimensional map (introduced by Kaplan and Yorke)

$$\begin{aligned}x_{n+1} &= ax_n \bmod 1 \\y_{n+1} &= by_n + \cos(2\pi x_n)\end{aligned}$$

for $a = 3$.

- (a) Show that the map has an attractor which lies in $|y| \leq 1/(1 - |b|)$ provided $|b| < 1$.
- (b) What is the Jacobean of the map at the point (x, y) ? What are the local directions of expansion and contraction?
- (c) What are the Lyapunov exponents of the map?
- (d) Now iterate the map numerically for $b = \frac{1}{4}$ and for $b = \frac{1}{2}$ e.g. using the program [2dmap](#) on the website (this map is labelled the “Yorke Map” there). Do the numerical values of the Lyapunov exponents agree with what you calculated?
- (e) Sketch the attractor of the map obtained numerically for $b = \frac{1}{4}$ and for $b = \frac{1}{2}$. Do you see any qualitative difference between the attractors at these two values?